

Heavy metal contamination in surface sediments of four important rivers leading to the Caspian Sea

Mohammad Abadi¹

Abbas Ali Zamani^{2*}

Abdolhossein Parizanganeh³

Younes Khosravi⁴

Hamid Badiie⁵

1, 2, 3. *Department of Environmental Science, Faculty of Sciences, University of Zanjan, Zanjan, Iran*

4, 5. *Department of Chemistry, Faculty of Science, Guilan University-University Campus, Rasht, Iran*

***Corresponding author:**

zamani@znu.ac.ir

Received date: 2018.01.21

Reception date: 2018.09.15

Abstract

Heavy metals tend to be adsorbed and accumulated in sediments as opposed to water or suspended solids. Sediments and biota bodies can act as a final sink for heavy metals in the aquatic environments. They accumulate through the food chain posing potential biological and ecological risks. Therefore, research on the heavy metal concentrations in aquatic ecosystems can help to protect human health and reduce the risks of heavy metal contamination. In this study, the contamination of heavy metals (Fe, Zn, Mn, Cu, Ni, Pb, Cd and Co) in surface sediments (0–5 cm) from 16 stations within four river intertidal zones (Neka, Tajan, Siahrood and Talar) were sampled and analyzed for metal contamination assessment in summer 2016. Results show that the mean concentration (mg kg^{-1}) of heavy metals in surface sediment samples followed the order; Fe (23300.00) > Zn (2028.87) > Mn (701.25) > Ni (36.77) > Pb (25.77) > Cu (19.62) > Co (18.62) > Cd (5.62). The mean concentration levels of the heavy metals Pb, Zn and Cd were more than mean background value. The statistical analyses showed that concentration of elements Fe, Mn, Co and Cu in the Siahrood River, Cd in Neka River and Zn in Tajan River is different ($P < 0.05$). Application of contamination factor (C_f), potential ecological risk factor of individual metals (E_r), enrichment factor (EF) and geo-accumulation index (I_{geo}) showed that the highest value of the above factors belongs to cadmium element. Also an assessment potential ecological risk index (RI), contamination degree (C_d) and pollution load index (PLI) suggests a considerable to very high ecological risk in Neka River. In addition to revealing potential ecological risks due to cadmium and other elements, our investigation markedly highlighted anthropogenic control over sediment quality deterioration and some immediate sediment quality management strategies are needed to remediate and control river bed contamination.

Keywords: Pollution, Environmental Indicators, Sediment, River, Caspian Sea.