Phylogenetic analysis study of Cultivable moderately halophit and halotolerant bacteria in Gomishan mud volcano

Mozhgan Ghiasian¹
Mohammad Ali Amoozegar^{2*}
Abbas Akhavan Sepahy³
Sara Saadatmand⁴
Mahmoud Shayandi⁵

- 1, 4. Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran
- 2. Extremophiles Laboratory,
 Department of
 Microbiology,Faculty of Biology,
 College of Sciences, University of
 Tehran, Tehran, Iran
- 3. Department of Microbiology, Islamic Azad University, North-Tehran Branch, Islamic Azad University, Tehran, Iran
- 5. Microbiology and Biotechnology Group, Research Institute of Petroleum Industry, Tehran, Iran

*Corresponding author: amoozegar@ut.ac.ir

Received date: 2017/10/20 Reception date: 2017/02/17

Abstract

Gomishan mud volcano is located 7 km from Gomishan city and sampling from this mud volcano was performed in August 2013. The temperature of mud was about 38 ± 1 °C with a pH of 8 ± 1 . The measured salinity was 34 g/L. Sodium and chloride were the highest and iron was the lowest ions of the mud. The culture medium for the isolation of moderately halophilic and halotolerant bacteria was used. High viable counts $(1-3\times10^6)$ were obtained in culture media. A total of 122 isolates were obtained 42 colonies were selected based on primarily morphological and physiological traits and their 16S rRNA sequences were determined. The isolated genera included Halomonas (20%), Arthrobacter (5%), Kocuria (5%), Thalassobacillus (5%), Marinobacter (20%), Paracoccus (5%), Roseovarius (5%), Jeotgalicoccus (5%), Bacillus (15%), and Staphylococcus (15%). Oxidase and catalase enzymes were studied in all isolates and 98.4% of isolates were positive catalase and 69.7% were positive oxidase.

Keywords: Mud volcano, Halotolerant bacteria, Moderately halophilic bacteria, Phylogenetic tree.