

## Effect of seasons on the composition and the ratio of omega-3 to omega-6 fatty acids of muscle tissue of *Barbus xantopterus*

Mehran Javaheri Baboli<sup>1\*</sup>

Mojhgan Khodadadi<sup>2</sup>

Mohsen Taghani niya<sup>3</sup>

1, 2. *Associated Professor, Department of Fisheries, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran*

3. *Graduated of Fisheries, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran*

**\*Corresponding author:**

mehranjavaheri@gmail.com

Received date: 2016.08.23

Reception date: 2017.04.26

### Abstract

The aim of this study was to determine the Seasonal variations in the muscle fatty acid composition of *Barbus xantopterus*. For this purpose, 21 samples of fish were caught in the middle month of each season during 2014-2015 and Total fatty acid compositions in 7 sample were investigated by a gas chromatographic method. The SFA content of the muscle were at the highest in the spring and significant difference with other season ( $P < 0.05$ ). The SFA content of the muscle were at the highest in the winter and were significantly reduced in the spring. Docosaehaenoic acid (22:6n-3), Eicosapentaenoic acid (20:5n-3), Linoleic acid (18:2n-6) and  $\gamma$ -linolenic (18:3n-6) were the most abundant PUFA. Omega-3 and docosaehaenoic fatty acids of the muscle tissue were not significantly different in different seasons. were dominant. Range of P/S ratio was from 0.52 to 1.36 in the different seasons. The changes range of IA and IT indices were 0.22 -0.96 and 0.25-0.66, respectively. In conclusion, it was shown that the saturated and mono unsaturated fatty acid compositions in the muscle of *B. Xantopterus* was significantly influenced by the seasons and poly unsaturated and mega3 fatty acids was not significantly influenced by the seasons.

**Keywords:** *Barbus xantopterus*, Muscle, Fatty acid Composition, Season, Shadegan Wetland.