The trend of changes in the Trohic State Index (TSI) in Chitgar Lake during $\gamma \cdot \gamma \gamma - \gamma \cdot \gamma q$

Ali Abedini^{`*} Siamak Bagheri^{`*} Ali reza Mirzajani[°] Ahmad Ghane['] Mohammad Reza Talakesh[°]

), r, r, ε. Inland water Aquaculture Research Center, Iranian Fisheries Science Research Institute, Agricultural Research Education Extension and (AREEO), **Organization** Bandar Anzali, Iran. o. Engineering Organization and Civil of Tehran city, Tehran, Iran

*Corresponding authors: aabedinim@yahoo.com

Received date: ۲۰۲۰,۰۰,۱۷ Reception date: ۲۰۲۰,۰۷,۱۳

Abstract

Lake of the Martyrs of the Persian Gulf (also known as the Chitgar Lake) is located in the north-western of Tehran. The construction of this artificial lake was done by Tehran Municipality was filled water of the Kan River in 7.17. TN, TP, SD and Chl-a was used to determine trophic levels. Water sampling was performed by Rotner sampler and the water quality was investigated on \circ stations site during $7 \cdot 17 \cdot 19$. Standard methods were used for Analysis of water parameters. To determine the difference between mean of the studied parameters, the One-way ANOVA variance analysis test and Duncan mean pair test was used, and the confidence level was 90%. Based on the study, the results indicated that during $7 \cdot 17 \cdot 19$ the mean yearly EC was 277 ± 10 μ s/cm, pH was $\forall, \forall 9 \pm \cdot, \epsilon \circ$ and SD was $\forall, \forall \pm \cdot, 9$ meter. The mean yearly of dissolve oxygen, total phosphorous, total nitrogen, respectively, were $\lambda, \pi_{\pm}, \xi, \cdot, \xi_{\pm}, \cdot, \xi_{\pm}, \cdot, \xi_{\pm}, \tau, \pi_{\pm}, \pi_{$ $\circ, \gamma_{\pm} \gamma, \gamma_{\mu} g/l$. During the months under study, the water temperature varied between $\circ, \Lambda_{\pm}, \sharp$ to $\Upsilon \Lambda, \P_{\pm}, \Upsilon$ as °C. The ratio of total nitrogen to total phosphorus was more than \mathcal{T} , which indicated phosphorus is limiting factor in the lake Eutrophication. Based on multivariate evaluation method, during $7 \cdot 17 \cdot 19$ mean of TSI was $\xi \cdot , 1 \pm 1, 7$ with range of $\gamma_{9-\circ}$. In the early years, TSI was oltraoligotrogh and during $(\cdot,)$ $(\cdot,)$ the eutrophication state close to the mesotrophic state and then it has almost returned to its original level during the downward trend after the summer of 7.1Å. Introducing fishes and refinery performance are two main factors that have contributed to the declining level of eutrophication in the lake.

Keywords: Water quality, Chitgar Lake, Physical and chemical parameter, TSI.