Assessment of concentrations of metal Cadmium and Cromium in different tissues of teal (Anas crecca)

Gholamreza Banagar^{1*} Hossein Alipour² Mehdi Hassanpour³

- 1. Department of Environmental, Bojnourd Branch, Islamic Azad University, Bojnourd, Iran.
- 2. Young Researchers and Elite Club, Bojnourd Branch, Islamic Azad University, Bojnourd, Iran.
- 3. Department of Environment, Provincial Directorate of Environment Protection, Golestan, Iran.
- *Corresponding author: gholam_banagar@yahoo.com

Received date: 2016/07/18 Reception date: 2017/09/21

Abstract

The aim of this study was to determine the concentration of metals Cd and Cr in liver, kidney and muscle tissues of teal Anas crecca (N=16) from Fereydoonkenar wetland. The study was conducted between November and December 2015. The elements Cd and Cr were assayed using a graphite furnace atomic absorption spectrometer and the results were given as mg/kg wet weight. The concentration of metals was compared between the liver, kidney and muscle tissue and using one-way ANOVA, followed by Tukey's test. The highest mean concentrations of Cd and Cr were observed in the kidney at 1.45 and 0.56 mg/kg wet weight, respectively. The lowest mean concentration of Cd and Cr were measured in muscle tissue at 0.05 and 0.08 mg/kg wet weight, respectively. The distribution patterns of Cd and Cr in tissues of teal follow the order: kidney>liver>muscle. All the results were statistically significant at P < 0.001. The results indicated that there were negative correlations (r = -0.798, P<0.01) between weight and Cr in muscle. In this study, Cd and Cr concentrations in the edible tissues of teal were below the maximum permissible limit of the WHO.

Keywords: Teal, Cd, Cr, Fereydoonkenar wetland.